



HPNA Position Statement

The Ethics of Use of Opioids in Palliative Nursing

It is the position of the Hospice and Palliative Nurses Association (HPNA) that:

- Hospice and palliative nurses and organizations must affirm that comprehensive and effective symptom management is a fundamental standard of care of all hospice and palliative care providers.
- Hospice and palliative nurses and organizations must affirm that symptom management should continue throughout the illness experience, including the final hours, days, and weeks of life.
- Hospice and palliative nurses and organizations must affirm that receiving safe and effective symptom relief is a right for all patients in every clinical setting, including older adults, infants, children, nonverbal, cognitively impaired patients, non-English-speaking people, and those with an active or a history of substance abuse.
- Hospice and palliative nurses must recognize that the risk of hastening death by administering opioids to patients with serious or life-threatening, progressive illnesses is minimal, particularly when administration occurs by or under the supervision of clinicians skilled in symptom management and is consistent with established management guidelines.
- Hospice and palliative nurses and organizations must recognize that the administration of opioids to alleviate suffering at the end of life is consistent with widely accepted ethical and legal principles.
- Hospice and palliative nurses and organizations must advocate for education across healthcare settings to help clinicians recognize their responsibility to relieve suffering at the end of life. This education should include information about the lack of empirical evidence proving opioids shorten life and the ethical and legal acceptability of administering opioids in doses sufficient to relieve symptoms.
- Healthcare organizations should develop policies regarding the administration of opioids for symptom relief at the end of life.
- Clinicians who regularly care for dying patients must achieve skill in providing aggressive symptom management.
- Hospice and palliative nurses and organizations must ensure access to opioids for those at the end of life.

Background

Quality palliative care starts with successful symptom management.^{1,2} While ensuring effective symptom management should be a primary objective in every clinical setting, it is especially important in those settings that provide end-of-life care.



Although experts agree that palliative care must focus on the prevention and relief of pain and suffering, clinicians, patients, and families may be reluctant to use opioids to achieve this goal, particularly at the end of life. This reluctance often stems from the fear that administering opioids may depress respirations and hasten death. However, there is no convincing scientific evidence that administering opioids using established guidelines, even in very high doses, accelerates death.³⁻⁵ Numerous clinical studies demonstrate no significant association among opioid use, respiratory depression, and shortened survival.⁶⁻⁹ Respiratory depression and other changes in breathing are part of the dying process and can more likely result from disease and multi-system organ failure than from opioids.^{5,10}

Most ethical concerns about causing harm, (i.e., hastening death) by administering opioids can be resolved through thoughtful application of established facts regarding the physiology of pain, the mechanism of drug action, and careful on-going assessment and reassessment of the patient's condition. Administering the specific drug or combination of drugs—no more and no less—required to relieve symptoms, using well validated treatment guidelines, is a fundamental responsibility of the nurse caring for any patient experiencing distressing symptoms.

Despite the lack of evidence that opioids hasten death, many clinicians continue to believe that administering opioids can accelerate the dying process.¹¹⁻¹³ For this reason, they seek moral justification for providing aggressive pain or symptom management using opioids. Per the principle of double effect—often used to justify interventions that have risk for harm—four conditions must be satisfied to establish a clinician's interventions to relieve pain and symptoms as morally permissible:

1. The act must be morally good or neutral, regardless of its consequences; relief of suffering by administering opioids is a priority in palliative care and therefore a "good act."
2. The clinician must intend the good effect (i.e., relief of the patient's suffering); although the bad effect (i.e., death) may be foreseen and permitted, it is not the clinician's intended effect.
3. The bad effect must not be the means by which the good effect is achieved; in other words, the patient does not need to die in order to be relieved of suffering.
4. The benefits of the good effect must outweigh the burdens of the bad effect; in this case, the benefits of achieving a reduction of suffering outweigh the minimal risk of hastening death.¹⁴

Today, although the principle of double effect may have academic merit, its application needs to be balanced with the growing body of evidence that using opioids appropriately is safe and effective and does not increase morbidity or hasten death. Nurses and other healthcare providers should not use the principle of double effect to perpetuate the myth that appropriate use of opioids hastens death.⁷ For some clinicians, this myth contributes to withholding adequate analgesia and under treatment of patient suffering.¹⁵



A broad consensus exists among professional groups, ethicists, courts, and many state legislatures that clinicians have a duty to administer opioids for symptom relief to patients at the end of life.^{2,16-18} Moreover, fear of hastening death because of opioid administration does not justify the withholding of pain medication.^{8,15-17}

Education and support are needed to ensure that clinicians in all settings understand their obligation to relieve pain and suffering and to achieve skill and confidence in the clinical activities that are necessary to meet this goal.^{16,18-22}

Definition of Terms

Pain: An unpleasant subjective sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage.²³

References

1. Hospice and Palliative Nurses Association. Pain Management at the End of Life. Pittsburgh, PA: Hospice and Palliative Nurses Association; 2017.
2. National Consensus Project for Quality Palliative Care. *Clinical Practice Guidelines for Quality Palliative Care*. 3rd ed. Pittsburgh, PA: National Consensus Project; 2013.
3. Portenoy RK, Sibirceva U, Smout R, et al. Opioid use and survival at the end of life: a survey of a hospice population. *J Pain Symptom Manage*. 2006;32(6):532-540. doi:10.1016/j.jpainsymman.2006.08.003.
4. Jennings AL, Davies AN, Higgins JP, Gibbs JS, Broadley KE. A systematic review of the use of opioids in the management of dyspnoea. *Thorax*. 2002;57(11):939-944.
5. Boland JW, Ziegler L, Boland EG, McDermid K, Bennett MI. Is regular systemic opioid analgesia associated with shorter survival in adult patients with cancer? A systematic literature review. *Pain*. 2015;156(11):2152- 2163. doi:10.1097/j.pain.0000000000000306.
6. Campbell ML, Bizek KS, Thill M. Patient responses during rapid terminal weaning from mechanical ventilation: a prospective study. *Crit Care Med*. 1999;27(1):73-77.
7. Sykes N, Thorns A. The use of opioids and sedatives at the end of life. *Lancet Oncol*. 2003;4(5):312-318.



8. Walsh TD, Rivera NI, Kaiko R. Oral morphine and respiratory function amongst hospice inpatients with advanced cancer. *Support Care Cancer*. 2003;11(12):780-784. doi:10.1007/s00520-003-0530-0.
9. Mercadante S, Villari P, Ferrera P, Mangione S, Casuccio A. The use of opioids for breakthrough pain in acute palliative care unit by using doses proportional to opioid basal regimen. *Clin J Pain*. 2010;26(4):306-309. doi:10.1097/AJP.0b013e3181c4458a.
10. Kamal AH, Maguire JM, Wheeler JL, Currow DC, Abernethy AP. Dyspnea review for the palliative care professional: assessment, burdens, and etiologies. *J Palliat Med*. 2011;14(10):1167-1172. doi:10.1089/jpm.2011.0109.
11. Solomon MZ, Sellers DE, Heller KS et al. New and lingering controversies in pediatric end-of-life care. *Pediatrics*. 2005;116(4):872-883. doi:10.1542/peds.2004-0905.
12. Bendiane MK, Bouhnik AD, Favre R, et al. Morphine prescription in end-of-life care and euthanasia: French home nurses' opinions. *J Opioid Manag*. 2007;3(1):21-26.
13. Bilsen J, Norup M, Deliens L, et al. Drugs used to alleviate symptoms with life shortening as a possible side effect: end-of-life care in six European countries. *J Pain Symptom Manage*. 2006;31(2):111-121. doi:10.1016/j.jpainsymman.2005.07.005.
14. Beauchamp TL, Childress JF. *Principles of Biomedical Ethics*. 7th ed. New York, NY: Oxford University Press; 2012.
15. Campbell ML. Treating distress at the end of life: the principle of double effect. *AACN Adv Crit Care*. 2008;19(3):340-344. doi:10.1097/01.AACN.0000330386.72260.96.
16. American Nurses Association. Registered Nurses' Roles and Responsibilities in Providing Expert Care and Counseling at the End of Life. Silver Spring, MD: American Nurses Association; 2010. <http://www.nursingworld.org/MainMenuCategories/Policy-Advocacy/Positions-and-Resolutions/ANAPositionStatements/Archives/etpain14426.pdf>. Accessed July 13, 2017.
17. Wash Rev Code ch 70, §122 (1992) apps.leg.wa.gov/RCW/default.aspx?cite=70.122&full=true. Accessed March 12, 2017.
18. Ferrell BR, Levy MH, Paice J. Managing pain from advanced cancer in the palliative care setting. *Clin J Oncol Nurs*. 2008;12(4):575-581. doi:10.1188/08.CJON.575-581.



19. Pierce SF, Dalton JA, Duffey M. The nurse's ethical obligation to relieve pain: actualizing the moral mandate. *J Nurs Law*. 2001;7(4):19-29.
20. Simone CB 2nd, Vapiwala N, Hampshire MK, Metz JM. Palliative care in the management of lung cancer: analgesic utilization and barriers to optimal pain management. *J Opioid Manag*. 2012;8(1):9-16.
21. Bookbinder M, McHugh M. Symptom management in palliative care and end of life care. *Nurs Clin North Am*. 2010;45(3):271-327. doi:10.1016/j.cnur.2010.04.002.
22. Mularski R, Puntillo K, Varkey B, et al. Pain management within the palliative and end-of-life care experience in the ICU. *Chest*. 2009;135(5):1360-1369. doi:10.1378/chest.08-2328.
23. American Pain Society. *Principles of Analgesic Use*. 7th ed. Glenview, IL: APS; 2016.

This statement reflects the best available evidence at the time of writing or revisions.

Approved by the HPNA Board of Directors April 2017

Copyright © 2017 by the Hospice and Palliative Nurses Association
To obtain copies of this statement please visit www.hpna.org